# BCG vaccine medical questionnaire (yoshin-hyo)(BCG 専用予診票)

<u>※医療機関の皆様へ:この予診票はみほんです。医師会への請求等は、日本語版の予診票を使用して下さい。</u> ※この予診票だけでは接種できません。こちらを参照に、正規の予診票(2枚複写)に転記して使用してください。 ※This english version yoshin-hyo cannot be use separately. Please transfers them to the Japanese version yoshin-hyo(2sheets).

(Eligible va	ccination age	: Befor	e reaching	age	1)
--------------	---------------	---------	------------	-----	----

Cannot be used at medical institutions outside of Yokohama Yokohama City Body temperature taken before examination:									
	Ward								
Address						Telephone No.			
Name of Person to									/
	Receive Vaccination					Date of		Age:	years
Parent/Guardian	Name of Parent/Guardian						(YY/MM/DD)		months
Turent/Guardian									Filled in
Questionnaire					Response Colun			in	by Physician
Have you read the Guide to Vaccinations for Children and other informational materials distributed by Yokohama City?					Yes		No		
	ur child's developmental histor	y:							
Birth weight:									
Was any abnormality d	letected at the time of delivery	?			Yes		No		
	letected following delivery?				Yes		No		
	letected during any of your chi		s an infant or young child?		Yes		1	No	
	hysically unwell in any way too	lay?			v	es	۲	No	
If so, please describe s	ymptoms in detail: (			)	1	65	1		
Has your child been ill	within the last month?				v	es	۲	No	
	nd date(s) of recovery: (			)	1	65	No		
Have any of the child's family members or other person around your child ever suffered (from birth until now) from tuberculosis?					Yes		1	No	
Has your child received any vaccination within the last month?					Yes		۲ I	No	
Type of vaccination a	<u>``</u>			)			1		
Has your child ever been diagnosed (from birth until now) with a congenital abnormality; a heart, kidney, liver and/or cranial nerve problem; immunodeficiency; or any other such condition?				Yes		1	No		
Disease/condition name(s): ( ) If applicable, did the doctor who examined your child for the above condition(s) approve of today's				Yes		1	No		
vaccination?									
Has your child every suffered a seizure (convulsions)?					Yes		1	No	
Approximate age: ( If so, did your child ha				)	Var		,	No	
	close relatives ever been diagn	and with concentral	immunadafiaianav?					No No	
			i minunodenciency?		Yes		1	NU	
Has your child ever felt unwell following vaccination?				)	Yes		1	No	
Type of vaccine: ( )   Has any of the child's close relatives ever felt unwell following vaccination?					Yes		No		
Do you have any questions about today's vaccination?								No	
To be filled out by the		1			1	63	1	10	
In light of the above q I have decided that too	juestionnaire responses and head day's vaccination will be (adm effects and side effects of the v	inistered / postpone					·		th, and other
effects and purpose of * Please circle either	camination and relevant explar f vaccination, the possibility of r "consent to" or "do not conse	severe side effects, ent to" in the parenth	relief measures in response theses above.	to adve	erse healt	h effects f	from vac	cination,	etc.
questionnaire form to	uestionnaire is to ensure safet Yokohama City.	y during vaccillation	n. Based on an understandin Parent/guardia	-			to the s		in or unis 「下さい
			-	-	_				
Lot no of Vaccine	Inoculum Dose	Admi	nistering Medical Institution	, Physi	ician's Na	ime and V	/accinati	ion Date	
	Subcutaneous inoculation with the								
	predetermined amount	Administering M	edical Institution:						
Lot no.	of vaccine using a	Administering Physician's Name: ※日本語版に記入して下さい					SIO -		
	tube-like apparatus containing multiple	Vaccination Date	: (YY/MM/DD): / /		ĸ ⊢ /≁ī				

needles

## ♦ ♦ Precautions Following BCG Vaccine Inoculation ♦ ♦

### Precautions Following Inoculation

- 1. Have the child monitored at the medical institution or ensure that a doctor can be contacted immediately for 30 minutes following the inoculation. Sudden side effects in reaction to the vaccination sometimes occur during this period.
- 2. Keep an eye out for the occurrence of side effects for four hours following inoculation.
- 3. Keep the injection sites clean. Although bathing is permitted following vaccination, <u>make sure</u> to avoid scrubbing or rubbing the injection sites.
- 4. If any unusual reaction occurs near the injection sites or the child's physical condition changes in any way, have the child examined by a doctor immediately.

#### Regular Reactions and Koch's Phenomenon

1. Regular Reactions

Red spots will appear **<u>about 10 days after inoculation</u>**, and they may contain small amounts of pus. This reaction is strongest about 4 weeks after injection but will heal naturally within 3 months' time. Avoid applying bandages and keep the area clean.

#### 2. Koch's Phenomenon

If the child has already been infected with tuberculosis, reddening, swelling, pus from holes created by the injection needles and other such symptoms may occur <u>within 10 days of</u> <u>inoculation</u>. This is known as Koch's phenomenon. If the injection sites are kept clean, these symptoms normally dissipate within 2–4 weeks.

However, if you observe a reaction that resembles Koch's phenomenon, your child must be examined to determine whether s/he has been infected with tuberculosis. Please contact the medical institution that administered the vaccine or your local ward office's Health and Welfare Center.

#### Side Effects of the BCH Vaccine

In rare cases, swelling of the lymph nodes in the armpit on the same arm as the injection sites may occur roughly one month after inoculation. If you keep an eye on their condition, they will return to normal. Visit a doctor if the area near the injection sites becomes inflamed, swells excessively, or accumulates and releases pus (after natural breakage).

#### Interval Until Next Inoculation

The next dose can be administered 27 days or more after the current BCG inoculation.